

On ε -Nets, Distance Oracles, and Metric Embeddings

Ilya Razenshteyn *

Abstract

We give two new applications of an observation from [ADF⁺11]. The first is an almost linear sized constant time data structure for reporting very large distances in undirected graphs. The second is a generic transformation of results about ℓ_1 -embeddability of metrics to a setting, where we are interested in preservation of large distances only.

1 Introduction

Let G be a weighted undirected graph with unique shortest paths (with non-negative weights). In [ADF⁺11] the following observation is heavily used: the set of all shortest paths of G (by shortest path we mean its set of vertices) has VC-dimension [VC71] at most two. We give two new applications of this fact.

1.1 Distance oracles

The first application is about distance oracles. Let $G = (V, E)$ be an undirected unweighted graph with n vertices. For vertices $v_1, v_2 \in V$ let $d(v_1, v_2)$ be the distance between v_1 and v_2 . Let ε, δ be two fixed positive constants. In Section 3 we build a data structure of size $O(n^{1+\delta})$ that given two vertices $v_1, v_2 \in V$ in constant time reports the following:

- “ $\perp$ ”, if $d(v_1, v_2) < \varepsilon n$;
- $d(v_1, v_2)$, if $d(v_1, v_2) \geq \varepsilon n$.

*Mathematics Department, Moscow State University, e-mail: ilyaraz@gmail.com

1.2 Metric embeddings

The second application is about metric embeddings. Let $G = (V, E, w)$ be an undirected weighted graph with n vertices. For vertices $v_1, v_2 \in V$ let $d(v_1, v_2)$ be the distance between v_1 and v_2 with respect to w , and $d_{\text{unw}}(v_1, v_2)$ be the distance with respect to unit weights.

Suppose we want to approximate metric d with ℓ_1 -norm. We are looking for mappings from V to ℓ_1 with small distortion.

Definition 1. Let us say that a mapping $f: V \rightarrow (\mathbb{R}^k, \ell_1)$ has distortion D if for every $v_1, v_2 \in V$

$$d(v_1, v_2) \leq \|f(v_1) - f(v_2)\|_1 \leq D \cdot d(v_1, v_2).$$

There are many results about ℓ_1 -embeddability. Let us state some of them.

Definition 2. Doubling dimension of d is the minimum integer k such that every subset $V' \subseteq V$ of diameter Δ can be covered with 2^k subsets of diameter $\Delta/2$.

Definition 3. Let us say that d is of negative type if $\sqrt{d}$ is isometrically embeddable into ℓ_2 .

Theorem 1. There are the following upper-bounds on D :

- [Bou85] For any d one can take $D = O(\log n)$;
- [Rao99] If G is H -minor free, then $D = O(\sqrt{\log n})$;
- [GKL03] If d has bounded doubling dimension, then $D = O(\sqrt{\log n})$;
- [ALN05] If d is of negative type, then $D = O(\sqrt{\log n} \log \log n)$.

But what if we are interested in preserving $d(v_1, v_2)$ only if $d_{\text{unw}}(v_1, v_2) \geq \varepsilon n$? In Section 4 we state and prove a generic black box transformation that allows us to replace all occurrences of n in Theorem 1 with $1/\varepsilon$ for this case.

In [ABC⁺05] somewhat stronger result about arbitrary, H -minor free, and bounded doubling dimension metrics is proved, but since our transformation is black box, it automatically holds for negative type metrics.

2 VC-dimension and ε -nets

Let X be a finite set. Let $R \subseteq 2^X$ be a system of subsets of X . The following definitions were given in [VC71].

Definition 4. Say that $U \subseteq X$ is shattered by R if for every $U' \subseteq U$ there exists $U'' \in R$ such that $U'' \cap U = U'$.

Definition 5. *VC-dimension of the pair (X, R) is the size of the largest subset of X , which is shattered by R .*

Consider the following set system: let $G = (V, E, w)$ be a weighted undirected graph with unique shortest paths. Let $X = V$, and $P \in R$ iff P is a shortest path in G . The following theorem was proved in [ADF⁺11]. We reprove it here for completeness.

Theorem 2. *VC-dimension of (X, R) is at most two.*

Proof. Let $\{v_1, v_2, v_3\} \subseteq V$. Suppose it is shattered by shortest paths in G . Then there exists a shortest path P that contains v_1, v_2 , and v_3 simultaneously. W.l.o.g. we can assume that v_1, v_2 , and v_3 appear in P exactly in the same order. But then, since shortest paths in G are unique, there is no shortest path that contains v_1 and v_3 , but does not contain v_2 . $\square$

If VC-dimension of (X, R) is small, then there exist small the so-called ε -nets.

Definition 6. *Let μ be a probabilistic measure over X , and ε be a fixed positive constant. Say that $N \subseteq X$ is an ε -net for (X, μ, R) if for every $U \in R$ such that $\mu(U) \geq \varepsilon$ the intersection of N and U is not empty.*

The following theorem was proved in [HW86].

Theorem 3. *If VC-dimension of (X, R) is at most d , then for every $\varepsilon > 0$ and every μ there exists an ε -net of (X, μ, R) of size $O(d \cdot \log(1/\varepsilon)/\varepsilon)$.*

By combining Theorem 2 and Theorem 3 we obtain the main ingredient of two our results.

Theorem 4. *Let $G = (V, E, w)$ be a weighted undirected graph with n vertices. Let d be a shortest-path metric on V with respect to w , and d_{unw} — with respect to unit weights.*

Then for every $\varepsilon > 0$ there exists a subset $U \subseteq V$ of size $O(\log(1/\varepsilon)/\varepsilon)$ such that for every $v_1, v_2 \in V$ such that $d_{\text{unw}}(v_1, v_2) \geq \varepsilon n$ there exists $u \in U$ such that $d(v_1, v_2) = d(v_1, u) + d(u, v_2)$.

Proof. By perturbing weights, we can assume that shortest paths in G are unique. Then we just apply Theorem 3 for uniform μ and $d = 2$ (by Theorem 2). $\square$

It follows from [Alo10] that one can not improve the bound on the size of U to $O(1/\varepsilon)$, but if G is H -minor free, then it follows from [KPR93] that there exists a desired U of size $O(1/\varepsilon)$.

3 Distance oracles

Let $G = (V, E)$ be an undirected unweighted graph. In this section we show how to build a small and effective distance oracle that reports large distances in G .

Theorem 5. *For every $\varepsilon > 0$ there exists an oracle of size $O(n)$ that given two vertices $v_1, v_2 \in V$ reports in constant time a number τ such that if $d(v_1, v_2) \geq \varepsilon n$, then $\tau = d(v_1, v_2)$.*

Proof. Let $U \subseteq V$ be a set from Theorem 4. Let us store for every vertex v distances to all elements of U . Then if we are given a query (v_1, v_2) , we report $\tau := \min_{u \in U} d(v_1, u) + d(u, v_2)$. By definition of U , if $d(v_1, v_2) \geq \varepsilon n$, then $\tau = d(v_1, v_2)$. $\square$

Since we want our oracle to distinguish the cases $d(v_1, v_2) < \varepsilon n$ and $d(v_1, v_2) \geq \varepsilon n$, we combine Theorem 5 with Thorup-Zwick oracle [TZ05].

Theorem 6 ([TZ05]). *For every positive integer k there exists a data structure of size $O(n^{1+1/k})$ that given two vertices $v_1, v_2 \in V$ reports in constant time a number D such that $d(v_1, v_2) \leq D \leq (2k - 1) \cdot d(v_1, v_2)$.*

Now we show how one can combine Theorem 5 and Theorem 6 and obtain the desired data structure.

Theorem 7. *For every $\varepsilon, \delta > 0$ there exists an oracle of size $O(n^{1+\delta})$ that given two vertices $v_1, v_2 \in V$ reports in constant time*

- “ $\perp$ ”, if $d(v_1, v_2) < \varepsilon n$;
- $d(v_1, v_2)$, if $d(v_1, v_2) \geq \varepsilon n$.

Proof. First, we build a Thorup-Zwick oracle for $k = \lceil 1/\delta \rceil$. Using it we can dismiss pairs (v_1, v_2) with $d(v_1, v_2) < \varepsilon n / (2k - 1)$. Now by using an oracle from Theorem 5 with $\varepsilon' = \varepsilon / (2k - 1)$ we can determine the exact value of $d(v_1, v_2)$. $\square$

4 Metric embeddings

In this section we state and prove a generic transformation of ℓ_1 -embeddability results of metric spaces to a setting, where we are concerned only about large distances.

Let $G = (V, E, w)$ be a weighted undirected graph with n vertices. Let d and d_{unw} be weighted and unweighted metric on V , respectively. Let $\varepsilon > 0$ be some parameter.

Definition 7. *Say that a mapping $f: V \rightarrow (\mathbb{R}^k, \ell_1)$ has ε -distortion D , if*

- For every $v_1, v_2 \in V$

$$\|f(v_1) - f(v_2)\|_1 \geq d(v_1, v_2);$$

- For every $v_1, v_2 \in V$ such that $d_{\text{unw}}(v_1, v_2) \geq \varepsilon n$

$$\|f(v_1) - f(v_2)\|_1 \leq D \cdot d(v_1, v_2).$$

Theorem 8. Suppose that for every $k \leq n$ and for every subset $U \subseteq V$ of size k there exists an embedding $g_U: U \rightarrow (\mathbb{R}^{t(k)}, \ell_1)$ with distortion $D(k)$.

Then for every $\varepsilon > 0$ there exists an embedding $f: V \rightarrow (\mathbb{R}^{t(O(\log(1/\varepsilon)/\varepsilon)) + O(\log n)}, \ell_1)$ with ε -distortion $O(D(O(\log(1/\varepsilon)/\varepsilon)))$.

Proof. Let $U \subseteq V$ be a set of size $k := O(\log(1/\varepsilon)/\varepsilon)$ from Theorem 4. There exists an embedding $g: U \rightarrow (\mathbb{R}^{t(k)}, \ell_1)$ with distortion $D(k)$.

For a vertex $v \in V$ let $p(v) \in U$ be the closest to v vertex from U (with respect to d). Consider the metric $d'(v_1, v_2) := d(v_1, p(v_1)) + d(v_2, p(v_2))$. It is well known that such metrics are embeddable into $(\mathbb{R}^{O(\log n)}, \ell_1)$ with distortion $O(1)$. Let $h: V \rightarrow (\mathbb{R}^{O(\log n)}, \ell_1)$ be a corresponding embedding.

Let us prove that a mapping $f: v \mapsto g(p(v)) \oplus h(v)$ has ε -distortion $O(D(k))$.

First, for every $v_1, v_2 \in V$

$$\begin{aligned} \|f(v_1) - f(v_2)\|_1 &= \|g(p(v_1)) - g(p(v_2))\|_1 + \|h(v_1) - h(v_2)\|_1 \geq \\ &\geq d(p(v_1), p(v_2)) + d'(v_1, v_2) = d(v_1, p(v_1)) + d(p(v_1), p(v_2)) + d(p(v_2), v_2) \geq \\ &\geq d(v_1, v_2). \end{aligned}$$

Second, suppose $d_{\text{unw}}(v_1, v_2) \geq \varepsilon n$. Then, by definition of U , there exists $u \in U$ such that $d(v_1, v_2) = d(v_1, u) + d(u, v_2)$. Thus,

$$\begin{aligned} \|f(v_1) - f(v_2)\|_1 &= \|g(p(v_1)) - g(p(v_2))\|_1 + \|h(v_1) - h(v_2)\|_1 \leq \\ &\leq D(k) \cdot d(p(v_1), p(v_2)) + O(1) \cdot d'(v_1, v_2) \leq \\ &\leq O(D(k)) \cdot (d(p(v_1), p(v_2)) + d(v_1, p(v_1)) + d(v_2, p(v_2))) \leq \\ &\leq O(D(k)) \cdot (d(v_1, v_2) + d(v_1, p(v_1)) + d(v_2, p(v_2))) \leq \\ &\leq O(D(k)) \cdot (d(v_1, v_2) + d(v_1, u) + d(v_2, u)) = O(D(k)) \cdot d(v_1, v_2). \end{aligned}$$

□

By combining Theorem 8 and Theorem 1 we obtain the following corollary.

Corollary 1. There are the following upper-bounds on ε -distortion D :

- For any d one can take $D = O(\log(1/\varepsilon))$;

- If G is H -minor free, then $D = O(\sqrt{\log(1/\epsilon)})$;
- If d has bounded doubling dimension, then $D = O(\sqrt{\log(1/\epsilon)})$;
- If d is of negative type, then $D = O(\sqrt{\log(1/\epsilon)} \log \log(1/\epsilon))$.

References

- [ABC⁺05] Ittai Abraham, Yair Bartal, T-H. Hubert Chan, Kedar Dhamdhere, Anupam Gupta, Jon Kleinberg, Ofer Neiman, and Aleksandrs Slivkins. Metric Embeddings with Relaxed Guarantees. *FOCS '05*, pages 83–100. IEEE Computer Society, 2005.
- [ADF⁺11] Ittai Abraham, Daniel Delling, Amos Fiat, Andrew Goldberg, and Renato Werneck. VC-Dimension and Shortest Path Algorithms. In *Automata, Languages and Programming*, volume 6755, pages 690–699. Springer Berlin / Heidelberg, 2011.
- [ALN05] Sanjeev Arora, James R. Lee, and Assaf Naor. Euclidean distortion and the sparsest cut. In *Proceedings of the thirty-seventh annual ACM symposium on Theory of computing*, STOC '05, pages 553–562. ACM, 2005.
- [Alo10] Noga Alon. A Non-linear Lower Bound for Planar Epsilon-Nets. In *Foundations of Computer Science (FOCS), 2010 51st Annual IEEE Symposium on*, pages 341–346, 2010.
- [Bou85] J. Bourgain. On lipschitz embedding of finite metric spaces in Hilbert space. *Israel Journal of Mathematics*, 52:46–52, 1985.
- [GKL03] Anupam Gupta, Robert Krauthgamer, and James R. Lee. Bounded geometries, fractals, and low-distortion embeddings. In *FOCS*, pages 534–543, 2003.
- [HW86] David Haussler and Emo Welzl. Epsilon-nets and simplex range queries. In *Proceedings of the second annual symposium on Computational geometry*, SCG '86, pages 61–71. ACM, 1986.
- [KPR93] Philip N. Klein, Serge A. Plotkin, and Satish Rao. Excluded minors, network decomposition, and multicommodity flow. In *STOC*, pages 682–690, 1993.
- [Rao99] Satish Rao. Small Distortion and Volume Preserving Embeddings for Planar and Euclidean Metrics. In *Symposium on Computational Geometry*, pages 300–306, 1999.

- [TZ05] Mikkell Thorup and Uri Zwick. Approximate distance oracles. *J. ACM*, 52(1):1–24, 2005.
- [VC71] V.N. Vapnik and A.Y. Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory of Probability and Its Applications*, 16(2):264–280, 1971.